




US 20210347783A1

(19) **United States**(12) **Patent Application Publication**
CHOI et al.(10) **Pub. No.: US 2021/0347783 A1**(43) **Pub. Date: Nov. 11, 2021**(54) **NOVEL DERIVATIVE OF
BETA-APOPICROPODOPHYLLIN AND
METHOD OF PREPARING THEREOF**(71) Applicant: **J&C Sciences**, Daejeon (KR)(72) Inventors: **Jong-Ryoo CHOI**, Daejeon (KR);
In-Young CHOI, Daejeon (KR);
So-Yeon NAM, Daejeon (KR); **Se Hee
HYUN**, Daejeon (KR)(21) Appl. No.: **16/991,190**(22) Filed: **Aug. 12, 2020**(30) **Foreign Application Priority Data**

May 11, 2020 (KR) 10-2020-0055887

Publication Classification(51) **Int. Cl.**
C07D 493/04 (2006.01)(52) **U.S. Cl.**
CPC **C07D 493/04** (2013.01)(57) **ABSTRACT**Provided are a novel derivative of β -apopicropodophyllin and a method of preparing the same, and more particularly,a compound represented by Formula 1 below, which is a novel derivative of β -apopicropodophyllin derived from podophyllotoxin, which is an anticancer agent, a method of preparing the same, and a composition for treating cancer, which includes the compound.

[Formula 1]

In Formula 1, R is a C_2 to C_{10} alkyl group, a C_2 to C_{10} alkyl group containing an allyl- or alkyne, a $-[CH_2]_n-C_3$ to C_8 cycloalkyl group, a substituted or unsubstituted $-[CH_2]_n$ -phenyl group, a substituted or unsubstituted $-[CH_2]_n-C_5$ to C_6 heteroaromatic group, a $-C(=O)-C_1$ to C_8 alkyl group, a substituted or unsubstituted $-C(=O)-[CH_2]_n$ -phenyl group, or a substituted or unsubstituted $-C(=O)-[CH_2]_n-C_5$ to C_6 heteroaromatic group, wherein n is an integer of 0 to 6.